

MODELS FOR THE OBSERVABLE SYSTEM PARAMETERS OF ULTRALUMINOUS X-RAY SOURCES

N. MADHUSUDHAN¹, S. RAPPAPORT¹, PH. PODSIADLOWSKI² & L. NELSON³

Draft version February 5, 2008

ABSTRACT

We investigate the evolution of the properties of model populations of ultraluminous X-ray sources (ULXs), consisting of a black-hole accretor in a binary with a donor star. We have computed models corresponding to three different populations of black-hole binaries, motivated by our previous studies. Two of the models invoke stellar-mass ($\sim 10 M_{\odot}$) black-hole binaries, generated with a binary population synthesis code, and the third model utilizes intermediate-mass ($\sim 1000 M_{\odot}$) black-hole accretors (IMBHs). For each of the three populations, we computed 30,000 binary evolution sequences using a full Henyey stellar evolution code. A scheme for calculating the optical flux from ULXs by including the reprocessed X-ray irradiation by, and the intrinsic viscous energy generation in, the accretion disk, as well as the optical flux from the donor star, is discussed. We present color-magnitude diagrams (CMDs) as “probability images” for the binaries as well as for the donor stars alone. “Probability images” in the plane of orbital period and X-ray luminosity are also computed. We show how a population of luminous X-ray sources in a cluster of stars evolves with time. The most probable ULX system parameters correspond to high-mass donors (of initial mass $\gtrsim 25 M_{\odot}$) with effective O through late B spectral types and equivalent luminosity classes of IV or V. We also find the most probable orbital periods of these systems to lie between 1-10 days. Estimates of the numbers of ULXs in a typical galaxy as a function of X-ray luminosity are also presented. From these studies we conclude that if the stellar-mass black-hole binaries are allowed to have super-Eddington limited X-ray luminosities: (i) the value of the binding energy parameter for the stellar envelope of the progenitor to the black-hole accretor must be in the range of $0.01 \lesssim \lambda \lesssim 0.03$ in order not to overproduce the ULXs, and (ii) the stellar-mass black-hole models still have a moderately difficult time explaining the observed ULX positions in the CMD. Other possible explanations for the apparent overproduction of very luminous X-ray sources in the case of stellar-mass black-hole accretors are discussed. Our model CMDs are compared with six ULX counterparts that have been discussed in the literature. The observed systems seem more closely related to model systems with very high-mass donors in binaries with IMBH accretors. We find that a significant contribution to the optical flux from the IMBH systems comes from *intrinsic* accretion disk radiation whose source is viscous dissipation of gravitational potential energy. In effect, the IMBH systems, when operating at their maximum luminosities ($10^{41} - 10^{42}$ ergs s⁻¹), are *milli-AGN*. With regard to the IMBH scenario, while attractive from the aspect of binary evolution models, it leaves open the larger question of how the IMBHs form, and how they capture massive stellar companions into just the correct orbits.

Subject headings: (stars:) binaries: general — X-rays: binaries — galaxies: star clusters — accretion, accretion disks — black hole physics

1. INTRODUCTION

An ultraluminous X-ray source (ULX) is an off-nucleus point-like source whose X-ray luminosity, L_x , exceeds 2×10^{39} ergs s⁻¹. An obvious candidate for such a source is an X-ray binary where a compact object accretes from a donor star. Under normal circumstances, the accretion rate is constrained by the Eddington limit of the accretor. The Eddington limit for a neutron star is $\sim 10^{38}$ ergs s⁻¹, and that for a $10 M_{\odot}$ black hole (BH) is $\sim 10^{39}$ ergs s⁻¹. Therefore, if the accretors are assumed to be stellar mass BHs, the observed luminosities of ULXs exceed the Eddington limit by factors of up to ~ 100 . The question as to how the Eddington limit could be violated remains a subject of considerable debate. On the other hand, if we assume the accretor to be an intermediate-mass black hole of $\sim 10^2 - 10^4 M_{\odot}$, the ULX luminosities can all be accounted for without violating the Eddington limit. If the accreting stars are indeed IMBHs, the question remains as to

how these objects are formed and how they acquire a massive stellar companion.

Several scenarios have been put forth to explain the apparent super-Eddington luminosities, assuming stellar-mass BHs. King et al. (2001) suggested that geometrical beaming in the direction of the observer, due to a thick accretion disk (Jaroszyński, Abramowicz, & Paczyński 1980), could lead to both a high luminosity and an even higher inferred luminosity. Körtling et al. (2002) make a similar argument with relativistic beaming, due to jets, as the explanation. In both of these scenarios, the emission is considered to be anisotropic. These somewhat contradict the observations of ionization nebulae around some ULXs (Pakull & Mirioni 2003) which tend to indicate the full implied isotropic luminosity. Another scenario proposed by Begelman (2002; 2006) suggests that photon bubble instabilities in the accretion disk can lead to an isotropic luminosity exceeding the Eddington limit by a factor of ~ 10 . Socrates & Davis (2006) invoke a hot, optically thin corona in conjunction with a geometrically thin, but optically thick, accretion disk to explain the observed ULX luminosities. Yet other scenarios invoke “slim” accretion disks to produce approximately isotropic luminosities that may exceed the Eddington limit by up to factors of ~ 10 (Abramowicz et

¹ Department of Physics and Kavli Institute for Astrophysics and Space Research, MIT, Cambridge, MA 02139; nmadhu@mit.edu

² Department of Astrophysics, Oxford University, Oxford OX1 3RH, UK

³ Physics Department, Bishop’s University, Sherbrooke, QC Canada J1M 0C8

al. 1988; Ebisawa et al. 2003). Although these scenarios have not been shown to work conclusively, they could, in principle, account for ULX luminosities up to $\sim 10^{40}$ ergs s $^{-1}$. In addition to these specific suggestions for bypassing the Eddington limit, the continuity and simplicity of the luminosity function of luminous X-ray sources from 10^{36} – 10^{40} ergs s $^{-1}$ (a simple power law; Grimm et al. 2003) has led some to conclude that these represent a single class of systems, i.e., neutron stars and stellar-mass black holes⁴ accreting from a normal donor star. However, we note that essentially none of these ideas involving stellar-mass BHs would plausibly be able to explain the ULXs at the higher end of the observed luminosities, namely those with $L_x \gtrsim 10^{40}$ ergs s $^{-1}$.

By contrast, the IMBH scenario, as suggested first by Colbert & Mushotzky (1999), accounts for the luminosities of all the ULXs because the Eddington limit for an IMBH in the mass range $\sim 10^2$ – $10^4 M_\odot$ varies between 10^{40} – 10^{42} ergs s $^{-1}$. Other evidence that may suggest the presence of an IMBH includes observations of mHz QPOs (Strohmayer & Mushotzky 2003) and ionization nebulae (Pakull & Mirioni 2003), and inferences of cool inner accretion disk temperatures (Miller et al. 2003; Miller et al. 2004; Cropper et al. 2004). Despite the several pieces of evidence supporting IMBH binaries as the model for ULXs, one of the main problems with this scenario lies in the unknown formation mechanism for such binaries. A few scenarios for IMBH formation have been proposed in the literature (see, for example, Portegies Zwart et al. 2004; Tutukov & Fedorova 2005) but a definitive scenario remains elusive. In particular, the evolution of supermassive stars with $M \gg 100 M_\odot$ is not well understood.

In three of our earlier papers, we studied the evolution of populations of binary systems pertaining to both models for ULXs, namely: (i) super-Eddington accretion onto LMBHs (Podsiadlowski et al. 2003, Rappaport et al. 2005); and, (ii) sub-Eddington accretion onto IMBHs (Madhusudhan et al. 2006). In those studies, the main objective was to investigate the evolution of the X-ray luminosities of the systems with time, and to estimate the formation efficiencies for each scenario. In Podsiadlowski et al. (2003) and Rappaport et al. (2005), we considered LMBHs in binary systems with donor stars in the mass range 2 – $17 M_\odot$. Some of the binary populations were generated using the binary population synthesis (BPS) code developed in Podsiadlowski et al. (2003). There, we showed that by allowing for a violation of the Eddington limit by a factor of ~ 10 – 30 , such systems would be able to explain most of the observed ULXs, except for the very most luminous ones. In Madhusudhan et al. (2006), we considered donor stars in binary systems with IMBHs with a representative mass of $1000 M_\odot$. There, we showed that in order to have a plausible formation efficiency for ULXs with $L_x \gtrsim 10^{40}$ ergs s $^{-1}$ the donor stars should be massive ($\gtrsim 8 M_\odot$) and the initial orbital separations should be close ($\lesssim 6$ – 40 times the radius of the donor star when on the main sequence).

A number of fairly secure optical identifications of ULX counterparts have now been reported in the literature. Liu et al. (2002) identified a unique counterpart to the ULX NGC 3031 X11 (in M81) and found it to be consistent with an O8 main sequence (MS) donor star. Kaaret et al. (2004) reported an optical counterpart to the ULX in Holmberg II to be con-

TABLE 1
MAGNITUDES AND COLORS OF ULX COUNTERPARTS
(ADAPTED FROM COPPERWHEAT ET AL. 2007)

System	M_B	M_V	$B - V$
NGC 4559 X-7 (C1)	-7.22 ± 0.19	-7.03 ± 0.16	-0.19 ± 0.25
M81 X-6	-4.28 ± 0.04	-4.18 ± 0.03	-0.10 ± 0.05
M101 ULX-1	-6.19 ± 0.15	-5.92 ± 0.12	-0.27 ± 0.19
NGC 5408 ULX	-6.40 ± 0.20	-6.40 ± 0.20	-0.00 ± 0.28
Holmberg II ULX	-6.03 ± 0.19	-5.78 ± 0.11	-0.25 ± 0.22
NGC 1313 X-2 (C1)	-4.70 ± 0.18	-4.50 ± 0.18	-0.20 ± 0.25

sistent with a donor star with spectral type between O4 V and B3 Ib. Kuntz et al. (2005) have studied the optical counterpart of M101 ULX-1 and found the colors to be consistent with those for a mid-B supergiant. Soria et al. (2005) studied candidate counterparts to the ULX in NGC 4559 and suggested the donor to be either a blue or a red supergiant of high mass (~ 10 – $20 M_\odot$). Mucciarelli et al. (2005; 2007) have found candidate optical counterparts to the ULX NGC 1313 X-2 that are either B0–O9 main-sequence stars or G supergiants.

The best candidate optical counterparts to the ULXs discussed above have been conveniently selected and summarized by Copperwheat et al. (2007). In particular, they tabulate the photometric values for the various systems, calculating reddening corrections and absolute magnitudes, wherever necessary. As can be seen from the above list, the observations seem to indicate that the spectral classes of the most promising candidates generally range between O and B spectral type, and the luminosity classes are either V or Ib. In the present study, we consider optical counterparts of six ULX systems for which photometric data in the B and V bands are available in the literature. The photometric data for these systems are given in Table 1 (as adapted from Copperwheat et al. 2007).

Theoretical studies of ULXs in the optical regime have heretofore concentrated on constraining the nature of the donor star, the mass of the accretor, and the orbital period. All the models follow the standard paradigm of ULXs being X-ray binaries with active accretion through Roche-lobe overflow from a donor star onto a BH accretor. Rappaport et al. (2005) discussed preliminary theoretical models involving disk irradiation in ULXs. They presented sample evolution tracks, on a color-magnitude diagram, for four ULX models, and discussed the optical appearance of ULX BH binaries. Pooley & Rappaport (2005) suggested detection of X-ray and/or optical eclipses as a means to constrain the mass of the accretor. In recent studies, Copperwheat et al. (2005; 2007) have also constructed irradiation models to describe the optical emission from ULXs, and used the models to constrain the properties of several systems observed in the optical. The parameters being considered were the mass, radius and age of the donor, and the BH mass in some cases. From the fits to the various observations, they find the counterparts to be consistent with being main-sequence stars or evolved giants/supergiants with spectral types O, B, or A.

In the present paper, we report a detailed population study spanning a large region of parameter space of ULX properties in an effort to better explore and help constrain ULX models. In particular, we investigate models of the optical properties of ULXs. We choose three sets of representative populations from our previous studies, and explore their optical proper-

⁴ In this paper, we refer to a stellar-mass black holes as a “low-mass black hole” or “LMBH”, rather than “SMBH” which might be confused with “supermassive black hole”.

ties and other observables. For the models with LMBHs, we choose two sets of populations obtained from the BPS code corresponding to two different prescriptions for the critical mass ratio in the primordial progenitor binary that determines whether a common envelope phase can occur. These two models yield very different distributions of secondary (i.e., ULX donor) star masses. For models with IMBHs, we choose model C from Madhusudhan et al. (2006). This model yielded the highest formation efficiency for ULXs in the IMBH scenario. For, each of the three populations, we follow 30,000 binary evolution calculations. The models are summarized in Table 2.

The X-ray luminosities are calculated using a standard formulation of Roche-lobe mass transfer in X-ray binaries. The optical flux from the system is determined as the sum-total of the optical flux from the donor star, X-ray irradiation of the accretion disk, and intrinsic energy generation in the disk. We present detailed color-magnitude diagrams (CMDs) for all the models. These include CMDs for the binary system, i.e., the sum of the optical flux of the donor and that due to radiation from the disk, as well as CMDs for the donor star alone. We also study in detail the evolution of the X-ray luminosity with the age of the system, t_{ev} , and the evolution of the X-ray luminosity with orbital period, P_{orb} .

2. METHODS

2.1. Binary Population Synthesis

In our previous studies we explored several models of ULX populations for both the low- (hereafter “LMBH”) and intermediate-mass (hereafter “IMBH”) black-hole scenarios (Rappaport et al. 2005; Madhusudhan et al. 2006). In the present work, we chose two LMBH- and one IMBH-binary population that were motivated by these previous studies. The parameters for the three models are summarized in Table 2, and the distributions of the initial system masses and orbital periods are shown in Figure 1. Models La and Lb are LMBH populations, whereas Model Ic represents an IMBH population.

To generate the LMBH binary populations, we used the binary population synthesis (BPS) code developed in Podsiadlowski et al. (2003). We briefly review here the formulation from Rappaport et al. (2005). We started with a very large set of massive primordial binaries and generated a much smaller subset of these that evolved to contain a black hole and a relatively unevolved companion star. The product was a set of “incipient” black-hole X-ray binaries with a particular distribution of orbital periods, P_{orb} , donor masses, M_{don} , and black-hole masses, M_{BH} , for each of a number of different sets of input assumptions (see, e.g., Fig. 2 of Podsiadlowski et al. 2003). For this part of the calculation, we employed various “prescriptions”, based on single-star evolution models for the primary, simple orbital dynamics associated with wind mass loss and transfer, assumptions about the magnitude of the wind mass loss from the primary as well as from the core of the primary after the common envelope, and natal kicks during the core collapse and formation of the black hole.

The decision in the binary population synthesis code of whether or not a common-envelope phase occurs, when the primary first overflows its Roche lobe and starts to transfer mass onto the secondary, is based on the evolutionary state of the primary when mass transfer commences. We first define R_{TAMS} and R_{HG} as the radii of the primary when it reaches the terminal main sequence and the end of the Hertzsprung

TABLE 2
ULX POPULATION MODELS

Model ^a	M_{BH}^b	M_{don}^c	$P_{\text{orb},i}^d$	λ^e	q_{crit}^f
La	6–15	5–18	1–18	0.1	1.2 or 2.0
Lb	6–15	5–27	1–12	0.1	1.2 → 2.0
Ic	1000	5–50	1–10	–	–

^aThe distributions of masses and periods for the different models are shown in Figure 1

^bApproximate range of black hole masses, in units of $M_{\odot}$

^cApproximate range of donor masses, in units of $M_{\odot}$

^dApproximate range of initial orbital periods, in units of days

^eDimensionless inverse binding energy of the envelope of the black-hole progenitor star

^fFor details of the prescription see eqns. (1) – (3) in the text.

gap, respectively. Given the initial orbital separation and the mass ratio of the primordial binary, $q \equiv M_{\text{prim}}/M_{\text{sec}}$, we can compute the Roche-lobe radius, R_L , of the primary. In order to form a relatively close black-hole binary of the type we are considering in this work (i.e., $P_{\text{orb}} \lesssim 20$ days), we require that the initial mass-transfer rate from the primary to the secondary be so large that it leads to a common-envelope and spiral-in phase. This depends on the initial mass ratio and evolutionary state of the secondary and is somewhat uncertain (see Pols 1994; Wellstein, Langer & Braun 2001). For one of our two models involving stellar-mass black hole accretors, we utilize the follow critical mass ratios, q_{crit} , depending on the relation among R_L , R_{TAMS} , and R_{HG} :

$$q_{\text{crit}} = 2 \quad \text{for } R_{\text{TAMS}} < R_L < R_{\text{HG}} \quad (1)$$

$$q_{\text{crit}} = 1.2 \quad \text{for } R_L > R_{\text{HG}} \quad (2)$$

For the second of our two models involving stellar-mass black-hole accretors we utilize an interpolated version of the above prescription:

$$q_{\text{crit}} = 2.0 - 0.8 \left[\frac{\log R_L - \log R_{\text{TAMS}}}{\log R_{\text{HG}} - \log R_{\text{TAMS}}} \right] \quad (3)$$

for $R_{\text{TAMS}} < R_L < R_{\text{HG}}$. This somewhat ad hoc prescription has the net effect of producing a much greater fraction of incipient ULX sources with high-mass donor stars.

Simple energetic arguments were used to yield the final-to-initial orbital separation during the common-envelope phase wherein the envelope of the primary is ejected. Here we utilized a parameter λ , which is the inverse of the binding energy of the primary envelope at the onset of the common-envelope phase in units of $GM_{\text{prim}}M_e/R_{\text{prim}}$, where M_{prim} , M_e and R_{prim} are the total mass, envelope mass, and radius of the primary, respectively. This parameter strongly affects the final orbital separation after the common-envelope phase, where smaller values of λ correspond to more tightly bound envelopes, and hence more compact post-common-envelope orbits.

With the above parameterization for the ejection of the common envelope we find the following expression for initial-final orbital separation:

$$\left(\frac{a_f}{a_i} \right)_{\text{CE}} \simeq \frac{M_c M_{\text{sec}}}{M_{\text{prim}}} \left(M_{\text{sec}} + \frac{2M_e}{\mathcal{E}_{\text{CE}} \lambda r_L} \right)^{-1}, \quad (4)$$

(e.g., Webbink 1985; Dewi & Tauris 2000; Pfahl et al. 2003), where the subscripts “prim”, “c”, and “e” stand for the progenitor of the black hole, its core, and its envelope, respectively, and “sec” is for the secondary, i.e., the progenitor of

the “donor star” in the black-hole binary. The quantity r_L is the Roche-lobe radius of the black-hole progenitor in units of a_i , $\mathcal{E}_{\text{CE}}$ is the fraction of the gravitational binding energy between the secondary and the core of the black-hole progenitor that is used to eject the common envelope, and λ is defined above. For typically adopted parameter values, $\lambda \sim 0.01 - 1$ (e.g., Dewi & Tauris 2000; Podsiadlowski et al. 2003), $\mathcal{E}_{\text{CE}} \simeq 1$, and $r_L \simeq 0.45 - 0.6$ (for an assumed mass ratio between the black-hole progenitor and the companion in the range of $\sim 2 : 1 \rightarrow 15 : 1$), the second term within the parentheses in eq. (4) dominates over the first. In this case, we find the following simplified expression for a_f/a_i :

$$\left(\frac{a_f}{a_i}\right)_{\text{CE}} \simeq \frac{r_L}{2} \left(\frac{M_c}{M_{\text{prim}} M_e}\right) M_{\text{sec}} \lambda \simeq 0.005 \left(\frac{M_{\text{sec}}}{M_{\odot}}\right) \lambda, \quad (5)$$

where the leading factor is $r_L/2 \simeq 1/4$, while the factor in parentheses involving the black-hole progenitor is $\sim 0.020 \pm 0.002 M_{\odot}^{-1}$ for virtually all of the progenitors we consider. This explains why the large majority of the incipient black-hole binaries (with low- to intermediate-mass donors) found by Podsiadlowski et al. (2003) resulted from an initially very wide orbit ($P_{\text{orb}} \sim \text{years}$ – when the primary attains radii of $\sim 1000 - 2300 R_{\odot}$) preceding the common-envelope phase in order to avoid a merger between the secondary and the core of the primary.

In the present work, for Models La and Lb (see Table 2 and Fig. 1), we chose two populations that were obtained using the BPS code described above. These two LMBH populations both utilize an inverse envelope binding energy parameter $\lambda = 0.1$, and two different prescriptions for the critical mass ratio required to produce a common envelope phase (see eqns. [1]–[3]). These result in quite different production rates for incipient ULXs with high-mass donor stars (i.e., $\gtrsim 15 M_{\odot}$).

For the IMBH case, Model Ic (see Table 2), we chose the same population as Model C in Madhusudhan et al. (2006). Due to our lack of a firm understanding of how IMBHs form in star clusters, and how they capture companion stars, we adopted a very simple prescription for the population of incipient IMBH binaries (but, see Blecha et al. 2006). For each binary, we chose the initial donor mass to lie uniformly in the range of $5 - 50 M_{\odot}$ using Monte Carlo methods. And, we chose the initial orbital separation uniformly over the range of 1–3 times the separation required for a ZAMS star of the corresponding mass to fill its Roche lobe. This translates to a range of initial orbital separations between $40 - 200 R_{\odot}$, and P_{orb} in the range of $\sim 1 - 10$ days. We found in our previous work that this population is the most favorable one we explored for producing ULXs in the IMBH scenario, i.e., the donor stars should be massive, i.e., $\gtrsim 8 M_{\odot}$ and the initial orbital separations, after circularization, should be close, $\sim 6 - 40$ times the radius of the donor star when on the ZAMS. This range of close orbital separations represents the case of direct tidal capture and circularization of a single field star by the IMBH (see, e.g., Hopman et al. 2004; Pfahl 2005).

For, each of the three modeled ULX populations, we carried out 30,000 binary evolution calculations. All models were run using 60 nodes of the `elix3` Beowulf cluster located at the University of Sherbrooke, Quebec. The run time for each of the three models was ~ 40 hours.

2.2. Stellar Evolution

The stellar evolution of the donor stars, including mass loss, was followed with `EZ` which is a stripped down, rewrit-




FIG. 1.— Distributions of initial donor masses, black hole masses and initial orbital periods of the LMBH systems. The red curves correspond to Model La and the green curves correspond to Model Lb. In the top panel, the solid curves represent donor masses and the dashed curves represent BH masses. For Model Ic, $M_{\text{BH}} = 1000 M_{\odot}$ and we choose M_{don} uniformly between $5 - 50 M_{\odot}$, and the initial orbital separation uniformly between 1-3 times the separation required for a ZAMS star of the corresponding donor mass to fill its Roche lobe.

ten version of a subset of the stellar evolution code developed by P. P. Eggleton (Paxton 2004). The physics of the program is unchanged from Eggleton’s (essentially as described in Pols et al. 1995), but the structure of the code has been modified to facilitate experiments involving programmed control of parameters. There are zero-age main-sequence (ZAMS) starting models for a variety of metallicities (from $Z = 10^{-4}$ to $Z = 0.03$) and masses (from 0.1 to $100 M_{\odot}$), with arbitrary starting masses created by interpolation. A user-provided procedure is called between steps of the evolution to inspect the current state, to make changes in parameters, and to decide when and what to record to log files. The source code and data for `EZ` can be downloaded from the web at <http://theory.kitp.ucsb.edu/~paxton>. For all models in this particular study, the number of stellar mesh-points was fixed at 200 in the interest of minimizing computation time.

2.3. Binary Evolution Calculations

The binary evolution was governed by a sequence of calculations involving the mass-transfer rate, the corresponding change in orbital separation, and the subsequent monitoring of mass loss. The mass transfer considered was due solely to Roche-lobe overflow. When the donor star fills its Roche lobe, the excess matter flows through the inner Lagrange point onto the accretor. Assuming spherical geometry, the mass-transfer rate was calculated using $\dot{M} \simeq 2\pi R H \rho v$, where R is the radius of the donor, H is the density scale height of the donor atmosphere, ρ is the density of the atmosphere at the Roche lobe, and v is the thermal velocity at the photosphere. Under the approximation of an isothermal, constant-gravity atmosphere, the scale height is given by $H \simeq kT/\mu g$, where μ is the mean molecular weight, and the density profile of the atmosphere is exponential. The Roche-lobe radius of the donor was taken to be $R_{\text{RL}} = 0.49 a q^{2/3} [0.6 q^{2/3} + \ln(1 + q^{1/3})]^{-1}$ (Eggleton 1983), where, a is the separation of the binary, $q = M_{\text{don}}/M_{\text{BH}}$, M_{don} is

the mass of the donor, and M_{BH} is the mass of the black hole. This procedure, while making use of some approximations, is self-adjusting to yield the correct mass-transfer rates.

The Eddington limited mass-transfer rate onto the accretor is given by $\dot{M}_{\text{Edd}} = 4\pi G M_{\text{acc}} / \eta \kappa c$, where G is the gravitational constant, κ is the radiative opacity, and η is the efficiency of the black-hole accretor in converting rest mass to radiant energy. The opacity is assumed to be predominantly due to electron scattering and is given by $\kappa = 0.2(1+X) \text{ cm}^2 \text{ g}^{-1}$, where, X is the hydrogen mass fraction. For a spinning black-hole, we take the accretion efficiency to be given by $\eta = 1 - \sqrt{1 - (M_0/3M_0)}$, where M_0 is the initial mass of the black-hole (Bardeen 1970). This assumes an initially non-spinning black-hole.

If we consider all the matter leaving the donor to be retained by the accretor, independent of the Eddington limit, then the resulting luminosity is referred to as the “potential luminosity” and is given by, $L_{\text{pot}} = \eta \dot{M} c^2$. The actual isotropic X-ray luminosity, on the other hand, is given by $L_x = \beta \eta \dot{M} c^2$, where β restricts the mass-transfer rate to the Eddington limit. For $\dot{M} < \dot{M}_{\text{Edd}}$, $\beta = 1$, otherwise $\beta = \dot{M}_{\text{Edd}}/\dot{M}$. When $\dot{M} > \dot{M}_{\text{Edd}}$, it is assumed that all the matter passes through the accretion disk until it reaches the inner edge of the disk, at which point the excess matter (above the Eddington limit) is ejected out of the system in the form of a jet. The mass and orbital angular momentum lost from the system in such ejection is incorporated when calculating the orbital separation of the system. We emphasize that we use the Eddington limited mass-transfer rate while calculating the orbital parameters during the evolution of the binary. However, along with the various orbital parameters, we also record L_{pot} at each step. And, in all the results presented in this paper, we typically refer to L_{pot} instead of the X-ray luminosity, L_x , of the system in order to examine the potentialities of violating the Eddington limit.

2.4. Disk Irradiation

The optical flux from a ULX is comprised of contributions from the donor star and from the reprocessing of X-ray photons by the accretion disk (as well as intrinsic energy generation within the disk). The irradiation of the accretion disk depends upon L_x , the geometrical properties of the disk, and whether or not the central accreting star has a hard surface. For purposes of the current calculation, we assume the disk to be geometrically thin. We find the effective temperature of the disk to be:

$$T(r) \simeq \left(\frac{L_x}{4\pi r_{\text{min}}^2 \sigma} \right)^{1/4} \left[\frac{4}{7} \xi'^2 x^{-10/7} (1-\alpha) + 3x^{-3} \right]^{1/4}. \quad (6)$$

Here, $x = r/r_{\text{min}}$, where r is the radial distance, r_{min} is the inner disk radius, taken to be $6GM_{\text{BH}}/c^2$, α is the X-ray albedo of the disk, which we take to be 0.7. The half thickness of the disk is given by $h(r) = \xi r^{9/7} r_{\text{max}}^{-2/7}$, and $\xi' = \xi (r_{\text{min}}/r_{\text{max}})^{2/7}$, where ξ is a constant equal to $h(r_{\text{max}})/r_{\text{max}}$ and r_{max} is the outer radius of the disk. We have taken r_{max} to be $0.7r_L a$ where r_L is the dimensionless Roche-lobe radius of the black-hole accretor. We somewhat arbitrarily adopt a value for ξ of 0.1, corresponding to a full angular thickness of the disk of $\sim 12^\circ$. The second term in the square brackets in eq. (6) comes from the Shakura-Sunyaev (1973) solutions for the temperature profile of a thin disk around a BH accretor. We have taken the inner edge of the accretion disk to be at the innermost stable circular orbit around a non-rotating BH. We

have neglected the factor $(1 - \sqrt{r_{\text{min}}/r})$ since its contribution to the optical flux can be assumed to be negligible. The first term in the square brackets in eq. (6) is adapted from the expression for the irradiation temperature of an accretion disk around a BH derived in King, Kolb & Szuszkiewicz (1997). This is a modified version of the corresponding term in eq. (8) of Rappaport et al. (2005) where we used an expression appropriate for radiation emanating from a centrally located, isotropically emitting, hard surface. The present formulation takes into account the fact that the X-ray radiation comes from the innermost part of the accretion disk surface and is emitted with a Lambertian angular distribution. Since the irradiated portion of the outer disk lies at a shallow angle with respect to the surface of the inner disk, the small value of the cosine factor significantly reduces the X-ray irradiation.

As an important caveat to our disk irradiation calculations, we note that the assumption of a thin disk (emitting in a Lambertian manner) likely breaks down at accretion rates slightly or greatly exceeding Eddington. In this case, the irradiation of the outer disk might become more or less efficient. However, such a calculation is considerably beyond the scope of the present work.

The optical flux from the disk is determined by using the temperature profile described in eq. (6), and assumes the local spectrum in each annulus of the disk is thermal. The radiant flux from each annular ring on the disk surface is determined from the specific intensity of the blackbody radiation and the temperature at that radius. The total flux due to the disk at a particular wavelength is calculated by integrating the annular flux over the radial extent of the disk, from r_{min} to r_{max} . The flux from the donor star is determined from its surface temperature, T_e , as obtained from the stellar evolution code. For the wavelengths under consideration, a thermal blackbody is a reasonable approximation in order to determine the flux from the star (with $\sim 10\%$ accuracy). This corresponds to an error in the apparent V magnitude of $\sim \pm 0.1$, which is within the error bars of most observations, and negligible compared to the absolute values of V and M_V . The corresponding error in $B-V$ is $\lesssim 0.2$.

The total flux of the system consists of contributions from the disk and the donor star. The B and V magnitudes were calculated from the total fluxes using $B = -2.5 \log(F_B/F_{B0})$ and $V = -2.5 \log(F_V/F_{V0})$, where F_B , F_{B0} , F_V and F_{V0} are the fluxes and reference fluxes at $\lambda = 4380 \text{ \AA}$ and $\lambda = 5450 \text{ \AA}$, respectively ($F_{B0} = 6.61 \times 10^{-9} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ \AA}^{-1}$ and $F_{V0} = 3.64 \times 10^{-9} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ \AA}^{-1}$).

After we compute $B-V$ for the combined radiation from the donor star and the accretion disk⁵, we apply a small correction ($\lesssim 0.2$ magnitudes) as a function of $B-V$ to take into account the treatment of the radiation as blackbody emission. These corrections were based on the differences between the $B-V$ colors computed from the simple algorithm given above and the colors of main sequence and supergiant stars tabulated by Johnson (1966).

2.5. Intrinsic Disk Emission

We have found from our models (see §3) that under certain circumstances the intrinsic disk emission (due to the viscous release of gravitational potential energy) which is represented by the second term in square brackets in eq. (6), can dominate the optical emission from the disk irradiation

⁵ All of our binary systems are assumed to be viewed at an average orbital inclination angle of 60° .




FIG. 2.— Illustrative temperature profiles of the irradiated accretion disk for an IMBH model (blue curves) and LMBH model (red curves). The solid curves are for the case of intrinsic (viscous) heating of the accretion disk alone, while the dashed curves are for the irradiation of the disk without any contributions from viscous heating. These correspond to the 2nd and the 1st terms, respectively, in the square brackets of eq. (6). The binary system parameters used for this illustrative example are given in the text.

(first term in square brackets). To illustrate this effect, we show in Fig. 2 the temperature profiles for an illustrative set of binary parameters: $L_{\text{pot}} = 10^{40}$ ergs s^{-1} , $P_{\text{orb}} = 30$ days, and $M_{\text{don}} = 10 M_{\odot}$. For an IMBH model ($M_{\text{BH}} = 1000 M_{\odot}$) the minimum and maximum disk radii are $r_{\text{min}} = 9 \times 10^8$ cm and $r_{\text{max}} = 1.5 \times 10^{13}$ cm, respectively, while for an LMBH model (with $M_{\text{BH}} = 10 M_{\odot}$), these radii are $r_{\text{min}} = 9 \times 10^6$ cm, and $r_{\text{max}} = 2 \times 10^{12}$ cm.

The red and blue curves in Fig. 2 are for the LMBH and IMBH models, respectively. The dashed curves are the temperature profiles for the case where irradiation is the only source of energy input to the disk; conversely, the solid curves are for the intrinsic (viscous) release of gravitational energy alone – without X-ray irradiation. For the LMBH model, we see that the intrinsic energy release dominates the irradiation contribution to the optical radiation only for radial distances in the disk of $\lesssim 1 R_{\odot}$. Since the optical light from the donor star itself comes from a typically much larger area, the intrinsic disk emission is generally not competitive with that from disk irradiation. By contrast the crossing point for the two temperature profiles corresponding to the IMBH model occurs at a radial distance of $\sim 55 R_{\odot}$ (and $T(r)$ remains $\gtrsim 10^4$ K out to $\gtrsim 20 R_{\odot}$). These radii are larger than the size of the donor star while it is near the main sequence or subgiant branch (i.e., while it is hot). Thus, it is quite possible that the largest contribution to the optical light from the IMBH systems comes from the intrinsic radiation emitted by the accretion disk itself. We will see the effect of this in §3 (see especially Fig. 6).

2.6. Irradiation of the Donor Star

We do not consider the effects of X-ray irradiation of the donor star. For our IMBH models the half angle subtended by the donor star ranges between $\sim 5^\circ$ and 10° . Since we have assumed an accretion disk with only a modest half angle of 6° , most or all of the donor star can be expected to be shielded by the accretion disk. For the LMBH models the donor stars, at

least initially, subtend larger half angles of $\sim 18^\circ$ to 24° . The fractional solid angle that the donors subtend ranges between ~ 0.025 and 0.036 while they are on the main sequence (and even less when they are on the giant branch and have transferred a significant amount of their mass). After taking into account shadowing by the accretion disk, the albedo of the donor star surface, and the Lambertian angular dependence of the disk irradiator, we find that no more than $0.0018 \rightarrow 0.0026$ of the X-ray luminosity is reprocessed on the face of the donor star. This is less than comes from the irradiated accretion disk and, furthermore, the larger distance between the X-ray source and the donor compared to that of the irradiated disk renders the flux even smaller yet. Therefore, for this work, we have neglected the irradiation of the donor stars.

3. RESULTS AND DISCUSSION

We present models of ultraluminous X-ray sources including calculations of the optical flux from the system. The optical flux includes contributions from the donor star as well as those due to radiation from the accretion disk. As mentioned previously, we have chosen three representative populations of binary systems for our study. Two of these populations have LMBHs and one population has IMBHs as accretors. At the beginning of each evolution, the donor star is on the zero age main sequence (ZAMS), and the initial system parameters are set by the binary population synthesis algorithms described above.

3.1. Evolution of a Single Binary

Figure 3 shows sample evolutionary tracks for three individual ULX binaries. The three binaries are illustrative of the three models listed in Table 2. Starting from the upper left panel, panel (a) shows the evolution of L_{pot} with the age of the system (t_{ev}). Panel (b) presents the evolution of L_{pot} with the orbital period, P_{orb} . Panel (c) shows the track in the color-magnitude diagram (CMD) of the ULX binary, taking into account the optical flux from the donor star as well as the flux due to radiation from the disk. And, panel (d) represents the CMD of the donor star alone. In calculating the optical flux from the disk, L_{pot} was used to irradiate the disk, which allows for possible violation of the Eddington limit. The evolution of the various parameters with time is denoted by the arrows. The blue curves represent the evolution of an IMBH binary with an initial donor mass, $M_{\text{don}} = 30 M_{\odot}$, a black-hole mass, $M_{\text{BH}} = 1000 M_{\odot}$, and an initial orbital period, $P_{\text{orb}} = 2.1$ days; taken from Model Ic (see Table 2). The green curves represent Model Lb with $M_{\text{don}} = 13 M_{\odot}$, $M_{\text{BH}} = 12 M_{\odot}$, and $P_{\text{orb}} = 1.9$ days. And, the red curves are for Model La with $M_{\text{don}} = 7 M_{\odot}$, $M_{\text{BH}} = 11 M_{\odot}$, and $P_{\text{orb}} = 1.6$ days. In order to illustrate the direction of evolution of the tracks, on each panel arrows are marked on the track that is most clearly separated from the others.

Let us consider the $L_{\text{pot}} - t_{\text{ev}}$ tracks in panel (a) for illustration, and follow the blue curve. The evolution starts with the donor on the ZAMS. As the donor evolves through the main sequence and somewhat beyond, the radius increases slightly and the donor fills its Roche lobe. Mass transfer then takes place onto the accretor through the inner Lagrange point. The $L_{\text{pot}} - t_{\text{ev}}$ curve depicts a modest increase in L_{pot} as the donor evolves through the main sequence. For high-mass stars, as are considered here, there is an overall contraction phase before the star ascends the giant branch. During the contraction phase, even a small decrease in radius leads to a large reduc-




FIG. 3.— Evolutionary tracks of illustrative binary evolution calculations. Panel (a): evolution of X-ray luminosity with age of the system. Panel (b): Orbital period versus X-ray luminosity. Panel (c): Color-magnitude diagram for the binary system. Panel (d): Color-magnitude diagram of the donor star alone. The blue curves show the evolution of an IMBH binary, typical of Model Ic, with an initial donor mass (M_{don}) of $30 M_{\odot}$, a black hole mass (M_{BH}) of $1000 M_{\odot}$, and an initial orbital period (P_{orb}), of 2.1 days. The green curves represent Model Lb with $M_{\text{don}} = 13 M_{\odot}$, $M_{\text{BH}} = 12 M_{\odot}$, and $P_{\text{orb}} = 1.9$ days. And, the red curves represent Model La with $M_{\text{don}} = 7 M_{\odot}$, $M_{\text{BH}} = 11 M_{\odot}$, and $P_{\text{orb}} = 1.6$ days.




FIG. 4.— Population diagrams for Model La, LMBH with $\lambda = 0.1$ and q_{crit} for the primordial binaries taking a value of 1.2 or 2.0 (see text, eqs. (1) and (2), and Table 2 for details). Panel (a): Evolution of potential X-ray luminosity with age of the system. Panel (b): Orbital period versus potential X-ray luminosity. Panel (c): Color-magnitude diagram of the binary systems. Panel (d): Color-magnitude diagram for donor stars alone. For both panels (c) and (d), contributions to the CMD image are made *only during times when* $L_{\text{pot}} \gtrsim 2 \times 10^{39} \text{ ergs s}^{-1}$. Each diagram is an image of 700×700 pixels, and represents the 30,000 X-ray binary evolution calculations that we computed. For each diagram, the parameter values from all the tracks were registered in each pixel that was traversed. The intensity in panel (a) represents the number of systems in the pixel, whereas the intensity in each of the other panels represents the accumulated evolution time spent in the pixel. The colors are scaled according to the 1/4 root of the corresponding intensities, with purple being of highest intensity and red being the lowest; the actual ratio of values between purple and red is $\sim 200:1$. In panel (c) the irradiation is taken to be from the full L_{pot} .

tion in the mass-transfer rate, $\dot{M}$, and hence the observed dip in L_{pot} . After the contraction phase, the star ascends the giant branch expanding through the Hertzsprung gap on a thermal timescale. This increase in radius leads to a spike in L_{pot} . The effects of the various phases of evolution of the donor star also manifest themselves through the other properties of the system. For example, the development of P_{orb} through the various evolutionary phases is evident from panel (b). One can see the sharp increase in the orbital period as the star expands

on the giant branch. The relatively smaller variations in L_{pot} as the period changes indicate the near constant, even if rapid, rate of growth of the donor star. Panels (c) and (d) depict the evolution of the optical properties of the system. As is apparent from panel (d), the increase in the absolute magnitude of the donor star as it ascends the giant branch takes place along with a corresponding decrease in its effective temperature. In the context of a binary system, however, some of the X-ray flux is reprocessed in the accretion disk and the tracks reflect




FIG. 5.— Population diagrams for Model Lb, LMBH with $\lambda = 0.1$ and q_{crit} for the primordial binaries taking a value over the range of 1.2–2.0 (see text, eq. (3), and Table 2 for details). Panel (a): Evolution of potential X-ray luminosity with age of the system. Panel (b): Orbital period vs. potential X-ray luminosity. Panel (c): Color-magnitude diagram of the binary systems. Panel (d): Color-magnitude diagram for donor stars alone. All other descriptors are the same as for Fig. 4.

the effects of mass transfer, as seen from panel (c).

3.2. Population Diagrams

Figures 4, 5 and 6 show color images of 30,000 evolutionary tracks each, corresponding to Models La, Lb and Ic, respectively (see Table 2). The panel arrangement and parameters shown are the same for all three figures, and match those of Fig. 3. Each panel contains a 700×700 image matrix of the corresponding parameters. For panel (a), the matrix covers 7 decades in L_{pot} and 3.5 decades in evolution time, in equally spaced logarithmic intervals. For each step of an evolutionary track, the position of the $L_{\text{pot}}-t_{\text{ev}}$ pair is located in the matrix and a value of ‘1’ is added to the corresponding matrix element. The resulting matrix, after recording all the

evolution steps from all the 30,000 tracks, is displayed as a color image, with 1/4-root scaling in intensity to enhance the dynamic range. For panel (b), the matrix covers 4 decades in P_{orb} and 7 decades in L_{pot} , in equally spaced logarithmic intervals. Each time the step of an evolution track lands in a matrix element, the evolution time-step is added to the corresponding matrix element, as opposed to adding a value of ‘1’ as was the case for panel (a). For panels (c) and (d), the matrices cover 11 units in the absolute visual magnitude, M_V , and 2 units in color index, in equally spaced linear intervals. Similar to the P_{orb} versus L_{pot} image in panel (b), a value equal to the evolution time step is added to the CMD matrix each time the step of an evolution track lands in a particular matrix element.




FIG. 6.— Population diagrams for IMBH Model Ic: Panel (a): Evolution of potential X-ray luminosity with age of the system. Panel (b): Orbital period versus potential X-ray luminosity. Panel (c): Color-magnitude diagram of the binary systems. Panel (d): Color-magnitude diagram for donor stars alone. All other descriptors are the same as for Fig. 4.

However, it is extremely important to note that contributions to the CMD are made *only during times when* $L_x \gtrsim 2 \times 10^{39} \text{ ergs s}^{-1}$ i.e., when the source would be a potential ULX.

As in panel (a), the matrices in the remaining three panels are also displayed as color images with 1/4-root scaling in intensity (i.e., relative probability) to enhance the dynamic range. The colors reflect this scaling, violet being of highest intensity and red being the lowest.

Thus, the intensity in panel (a) is a measure of the number of systems with a particular L_{pot} found at a particular time in the evolution of the cluster. These numbers allow one to calculate the number of active ULX systems in the population at any given epoch and to make estimates of the numbers of

ULXs of a certain L_{pot} in typical galaxies. For all the other images, the intensity at a point is a measure of the total duration (time) spent by all the systems in that interval of parameter space. Since this intensity incorporates both the numbers of systems and the amount of time each system spends in that interval of parameter space, it provides a probability map of the corresponding parameter space. For purposes of discussion later, we shall refer to violet and dark blue regions in these diagrams as regions of high probability and to regions in red as those of low probability.

3.3. Evolutionary Phases of the Donor Stars

Comparing panel (c) from Fig. 3, to that of Figs. 4, 5, and 6, we see that the regions of high probability correspond to

the early phases of the donor star's evolution, i.e., on the main sequence or the sub-giant branch phase. This is apparent because the donor star spends a predominant amount of its lifetime on the main sequence, and evolves rather quickly on the giant branch. Consequently, the regions of low probability on the diagram are the regions corresponding to the giant-branch phase of the evolutionary tracks or other regions not readily accessed during the binary evolution. However, this description of the phases of evolution with respect to regions on the CMD is only approximate. The exact structure of the CMD is a result of a complex interplay between the evolution of the donor star, its mass loss, and the radiation from the disk at a particular evolutionary stage. From panel (c) of Figs. 4, 5, and 6, we find that, with a high probability, the donor stars belong to the equivalent luminosity classes for single stars of IV and V, with a $B-V$ color range of about -0.35 to -0.10 (corresponding to an effective spectral class of O through late B).

In many of the panels in Figs. 4, 5, and 6 one can discern some of the individual tracks, roughly parallel to each other in the initial stages of the evolution. Tracks higher on the diagram (i.e., lower M_V) correspond to higher-mass donors since, at any evolution phase, more massive stars are brighter. Higher-mass donor stars become progressively rarer and also spend lesser amounts of time on the main sequence; hence the probability fades toward lower M_V .

3.4. Relevance to the Observations

The data from the optical observations of six ULXs discussed earlier (see Table 1) are plotted along with the model CMDs. For models La (Fig. 4) and Lb (Fig. 5), we see that two of the six data points fall on high-probability regions of the CMD (panel c), two on regions of moderate probability, and two on low-probability regions. However, for the IMBH scenario all six data points fall on or close to the high-probability regions of the CMD in Fig. 6. Thus, in the framework of the models presented here, the IMBH scenario appears to be the more favorable one.

We also see from Fig. 6 that all the observational points fall in regions corresponding to the very high donor masses. This indicates how the IMBH models fit the data better than the LMBH models. The brighter magnitudes of the observed systems require massive donor stars. And, massive donor stars need more massive accretors in order for mass transfer to occur stably via Roche-lobe overflow. In the LMBH scenario, the donor masses are limited by the fact that they cannot be more than about twice the mass of the BH accretor in order for stable mass transfer to take place. In practice, this limits most of the donor masses to be $\lesssim 20 M_\odot$. In the IMBH case, on the other hand, the BH mass has been set at $1000 M_\odot$ and the maximum donor mass we consider is $50 M_\odot$. Quite clearly, it is the higher-mass donors ($M \gtrsim 25 M_\odot$) that lead to the high probability region where the observed systems lie. Such high-mass donors in the LMBH models are relatively scarce.

3.5. Effects of Radiation from the Accretion Disk

Panel (d) of each figure shows the CMD of the donor stars alone. Comparing panels (c) and (d) in each figure, we see that the difference is most apparent in the IMBH scenario. Comparing panels (c) and (d) in Fig. 4, we see that the CMDs differ significantly only in regions corresponding to the giant-branch phases of the evolutionary tracks, and remain quite

similar over the rest of the diagrams. This observation is similar for Fig. 5. And since the giant-branch phases lie in the low probability regions of the diagram, one can say that the optical flux from disk radiation contributes relatively little to the observed optical appearances of ULXs in the LMBH scenario. By contrast, for the IMBH model, a comparison of panels (c) and (d) in Fig. 6 shows that the contribution of disk irradiation and *intrinsic* disk radiation from viscous losses (see §2.5) is quite significant in this case, and has substantially altered the CMDs in most phases of the evolution.

The greater influences of the disk radiation for the IMBH model (see Fig. 6) compared to those for the LMBH models seen in Figs. 4 and 5 can be explained by two effects. First, in general, the IMBH systems have higher luminosities by about an order of magnitude – largely by virtue of their higher mass donor stars. This obviously increases the power going into disk irradiation. However, a comparison of IMBH and LMBH systems at the *same* values of P_{orb} and L_{pot} reveals an important difference. The disks in IMBH systems still emit much more optical power than do the LMBH models, and this is predominantly at the blue end of the spectrum. The reason is that there is a substantial amount of intrinsic radiation from the disk (viscous release of gravitational potential energy; see §2.5). The physical explanation for this is straightforward. At the same radial distance from the central black hole, the IMBH system releases ~ 100 times more gravitational potential energy than the corresponding LMBH system (simply due to the higher mass of the black hole). This release can end up dominating over the disk irradiation in regions where the bulk of the optical emission is released (see Fig. 2).

Finally, it should be noted that the CMD for the donor star in a binary is expected to be different from that of a single star because of the mass loss via Roche-lobe overflow.

3.6. Orbital Period versus X-Ray Luminosity

Panel (b) in each of Figures 3, 4, 5, and 6 shows the evolution of L_{pot} with P_{orb} . As can be seen in Fig. 3, the variations in P_{orb} reflect the changes in radius of the donor star through the various phases of evolution. Owing to the dependence of the Roche-lobe radius on the orbital separation, any rapid increase in the radius of the star while in contact with its Roche lobe causes a rapid increase in the orbital separation, and hence in P_{orb} . The final rapid rise in P_{orb} at the highest L_{pot} characterizes the expansion of the donor on the giant branch, leading to long orbital periods and high mass-transfer rates. As is apparent, the long values of P_{orb} in the different models indicate the giant-branch phases of the donors, and the short period regions pertain to the main-sequence and sub-giant phases of evolution. The intensity at a given location in panel (b) is proportional to the number of systems passing through the region and the amount of time spent by the systems in that region. Consequently, the intensity in that region is a measure of the probability of finding a system at that location in parameter space. We then see that, for any model, the most probable regions for $L_{\text{pot}} \gtrsim 10^{39}$ ergs s $^{-1}$ lie within a period range of ~ 1 -10 days.

It follows that the total probability of a system to be in the giant-branch phase is proportional to the sum of all the evolution times recorded in the long-period region, taken to be $P_{\text{orb}} \gtrsim 10$ days. Similarly, the total probability of being in the pre-giant phase is proportional to the sum of all the evolution times recorded in the short-period region, i.e., $P_{\text{orb}} \lesssim 10$ days. Thus, we can calculate the relative probability for a ULX donor to be in the giant-branch phase versus that to be




FIG. 7. — The fraction of LMBH/IMBH binaries as a function of star cluster evolution time for three different lower limits on the X-ray luminosity. The different models are labeled in the panels. For each model, the blue curves represent fractions of the initial systems that have $L_{\text{pot}} \gtrsim 10^{36}$ ergs s $^{-1}$, the green curves represent fractions with $L_{\text{pot}} \gtrsim 2 \times 10^{39}$ ergs s $^{-1}$, i.e., ULX luminosities, and the red curves are for fractions of systems with $L_{\text{pot}} \gtrsim 10^{40}$ ergs s $^{-1}$.

on the pre-giant branch by calculating the ratios of the two sums for all $L_{\text{pot}} \gtrsim 2 \times 10^{39}$ ergs s $^{-1}$. The relative probabilities thus calculated are 0.08, 0.05 and 0.15 for Models La, Lb and Lc respectively. This reflects, in a quantitative way, the known fact that the systems are more likely to be found in the pre-giant phase than in the giant phase because of the relatively small amount of time they spend on the giant branch.

The implication of the $P_{\text{orb}}-L_{\text{pot}}$ population diagram lies in the fact that given an observed value of L_{pot} , one can estimate the most probable orbital period corresponding to each of the three ULX scenarios presented in Figs. 4, 5 and 6. For instance, given an observed $L_{\text{pot}} = 10^{40}$ ergs s $^{-1}$, we see from Figs. 4, 5 and 6 that the most probable periods lie between 1–5 days, 1–5 days, and 2–10 days for the Models La, Lb and Lc respectively. Thus, the diagrams serve as rough guidelines showing what orbital periods are to be expected.

3.7. Evolution of the X-Ray Luminosity

Panel (a) of each of Figs. 4, 5 and 6 shows a population diagram of the evolution of the X-ray luminosity (L_{pot}) with the age of the system. Defining the ULX luminosities as $L_{\text{pot}} \gtrsim 2 \times 10^{39}$ ergs s $^{-1}$, we see that the systems in Model Lc are most active as ULX systems between ~ 2 and 15 Myr while systems in Models La and Lb are most active between ~ 4 and 25 Myr. These modest differences in active cluster lifetimes can be explained on the basis of the ranges of donor masses in the different models. Donors with higher masses have shorter nuclear evolution timescales, and hence shorter overall lifetimes. So, the higher the donor masses, the shorter will be the active lifetimes of the systems, leading to shorter overall timescales over which a star cluster containing such binaries is X-ray active. Shorter lifetimes of systems also mean that the mass-transfer rates are higher, leading to systems with higher luminosities. Consequently, while systems in Model Lc can produce high L_{pot} systems even before as-

cending the giant branch, systems with lower masses have to wait longer to produce their peak X-ray luminosities. It also follows that the numbers of systems having high L_{pot} will contain a higher proportion of higher-mass donors.

Figure 7 elucidates this latter point. It shows plots of fractions of the initial systems that have L_{pot} greater than 10^{36} , 2×10^{39} and 10^{40} ergs s $^{-1}$, at any given evolution time. We see that the maximum percentage of systems having (potential) ULX luminosities at any time during the age of a star cluster is $\sim 20\%$ for Model La, $\sim 30\%$ for Model Lb, and $\sim 25\%$ for Model Lc. We also see that the percentages of systems having $L_{\text{pot}} > 10^{40}$ ergs s $^{-1}$ are $\sim 2\%$, $\sim 3\%$, and $\sim 20\%$, respectively, for these same models. It follows that the IMBH model allows for the largest numbers of very high luminosity systems. The percentages quoted above refer only to the most active phase of the corresponding clusters. The detailed evolution of the percentages with the star cluster age can be seen from the figure. However, the very small numbers of systems with $L_{\text{pot}} > 10^{40}$ ergs s $^{-1}$ at *any* time during the entire cluster lifetime make the LMBH models an unlikely scenario for the most luminous ULXs (independent of Eddington-limit arguments).

3.8. Estimates of ULX Numbers

Based on the $L_{\text{pot}}-t_{\text{ev}}$ population diagrams, we can also make rough estimates of the numbers of ULXs of different L_{pot} that can be found in a galaxy in *steady state*, assuming the different models of ULX formation.

Each element of the $L_{\text{pot}}-t_{\text{ev}}$ matrix contains the number of BH binary systems that have luminosities between L_{pot} and $L_{\text{pot}} + \Delta L_{\text{pot}}$ and ages between t_{ev} and $t_{\text{ev}} + \Delta t_{\text{ev}}$. Both, L_{pot} and t_{ev} are logarithmically spaced with 100 bins per decade. Given the fact that we evolve 30,000 binary systems for each model, the fraction of all high-mass binary systems that lie in the above mentioned intervals of L_{pot} and t_{ev} is given by $m(L_{\text{pot}}, t_{\text{ev}})/30,000$, where m denotes the 700×700 $L_{\text{pot}}-t_{\text{ev}}$ matrix.

For the LMBH models, the BH accretors in this age interval, Δt_{ev} , have all resulted from core-collapse supernovae (SNe) in an equal time interval at some epoch, t_{ev} , in the past. Assuming a uniform SN core-collapse rate of R_{SN} , the total number of BH binaries formed in this time interval is given by $R_{\text{SN}} \times f_{\text{BH}} \times \Delta t_{\text{ev}}$, where f_{BH} is the fraction of all core-collapse supernovae that result in BH binaries with high mass donors (as determined from the BPS code). It then follows that the number of binary systems with luminosities between L_{pot} and $L_{\text{pot}} + \Delta L_{\text{pot}}$ and evolutionary ages between t_{ev} and $t_{\text{ev}} + \Delta t_{\text{ev}}$ is given by:

$$\Delta N(L_{\text{pot},j}, t_{\text{ev},i}) = \frac{R_{\text{SN}} \times f_{\text{BH}}}{30,000} \times \Delta t_{\text{ev},i} \times m(L_{\text{pot},j}, t_{\text{ev},i}), \quad (7)$$

where j and i label the matrix element bins of L_{pot} and t_{ev} , respectively. The total number of sources with $L_x > L_{\text{pot}}$ in a galaxy in steady-state can be obtained by summing over all the time bins of t_{ev} , and all the luminosity bins greater than L_{pot} , as:

$$N(> L_{\text{pot},k}) = \frac{R_{\text{SN}} \times f_{\text{BH}}}{30,000} \times \sum_{j=k}^{700} \sum_{i=1}^{700} \Delta t_{\text{ev},i} \times m(L_{\text{pot},j}, t_{\text{ev},i}). \quad (8)$$

For illustration, we choose a uniform supernova rate $R_{\text{SN}} = 0.01$ yr $^{-1}$ (e.g., to represent a typical Milky-Way type galaxy),




FIG. 8.— The expected numbers of ULXs at the present epoch as a function of X-ray luminosity in a starburst galaxy with a core collapse supernova rate of 0.01 per year. The red, green, and blue curves correspond to Models La, Lb and Ic, respectively.

and $f_{\text{BH}} \simeq 3.75 \times 10^{-4}$ and 9.44×10^{-4} for models La and Lb, respectively. These values of f_{BH} are deduced from the results of our BPS code. The computed steady-state cumulative luminosity functions for Models La and Lb are shown in the red and green curves of Fig. 8, respectively.

For the IMBH model, we consider the IMBHs to be formed dynamically in young star clusters. Following a formulation similar to eq. (8) for LMBH models, we find for the IMBH model:

$$N(> L_{\text{pot},k}) = \frac{R_{\text{ysc}} \times f_{\text{IMBH}} \times f_{\text{cap}}}{30,000} \times \sum_{j=k}^{700} \sum_{i=1}^{700} \Delta t_{\text{ev},i} \times m(L_{\text{pot},j}, t_{\text{ev},i}), \quad (9)$$

where, R_{ysc} is the formation rate of young star clusters (with $M_{\text{clus}} \gtrsim 10^4 M_{\odot}$) in a typical spiral galaxy, f_{IMBH} is the fraction of all such clusters that form an IMBH and f_{cap} is the fraction of all such IMBHs that capture a massive companion into a close orbit. As illustrative values, we choose a uniform $R_{\text{ysc}} = 10^{-5} \text{ yr}^{-1}$ (estimated from the combined work of de Grijs et al. 2003; Hunter et al. 2003; Grimm et al. 2003), $f_{\text{IMBH}} \simeq 0.1$, and $f_{\text{cap}} \simeq 0.05$ (see, e.g., Blecha et al. 2006). The distribution of the numbers of systems as a function of L_{pot} is shown by the blue curve in Fig. 8.

It can be seen from Fig. 8 that the expected numbers of ULX systems per galaxy are in fair agreement with the observations (Ptak & Colbert 2004; Grimm et al. 2003) for the IMBH model, albeit with very large uncertainties. Specifically, these estimates in Fig. 8 indicate that a typical galaxy is expected to harbor ~ 0.07 and ~ 0.03 ULXs with $L_{\text{pot}} > 2 \times 10^{39}$ and $10^{40} \text{ ergs s}^{-1}$, respectively. By contrast, for the LMBH models, the expected numbers range from about ~ 12 – 20 for $L_{\text{pot}} > 2 \times 10^{39} \text{ ergs s}^{-1}$ per galaxy to ~ 1.4 – 2.7 with $L_{\text{pot}} > 10^{40} \text{ ergs s}^{-1}$. These numbers seem to be too large by about a factor of ~ 100 compared with the observations (Ptak & Colbert 2004; Grimm et al. 2003). Moreover, there is considerably less flexibility, or uncertainty, in the LMBH models in comparison with the IMBH models. And, while one might dismiss the number of predicted ULXs with $L_{\text{pot}} > 10^{40} \text{ ergs s}^{-1}$ as violating the Eddington limit beyond the realm of the plausible, the predicted systems with $L_{\text{pot}} > 2 \times 10^{39} \text{ ergs s}^{-1}$ are not so easily dismissed. One completely straightforward

way of reducing the computed numbers of ULX systems in the LMBH model is to adjust the λ parameter downward (recall that λ is the dimensionless inverse binding energy of the stellar envelope of the BH progenitor). From the output of our BPS code, and an examination of Fig. 3 in Podsiadlowski et al. 2003, we can see that a value of λ in the range 0.01 – 0.03 will decrease the rate of production of BH binaries with high-mass donor stars by factors of ~ 10 – 100 . Such values of λ are quite consistent with what stellar structure calculations tell us (see Fig. 1 of Podsiadlowski et al 2003; Dewi & Tauris 2000).

Overall, there are a number of uncertainties associated with the various parameter values that we have used to calculate the ULX number estimates from eqns. (8) and (9). The primary utility of Fig. 8 beyond providing crude estimates is that, given more accurate values for the uncertain input parameters, one can find the corresponding expected numbers of ULXs by simply scaling the expected numbers inferred from the figure.

4. SUMMARY AND CONCLUSIONS

In this paper, we have explored the optical properties and other observable parameters for a range of ULX models. We investigated a very large number of systems from three different ULX populations, two consisting of stellar-mass BH accretors, which we refer to as stellar-mass (i.e., “low-mass”) black holes (LMBHs), and one consisting of intermediate-mass black-hole accretors (IMBHs). For each population, we computed the evolution of 30,000 individual binary systems, and generated population diagrams to explore a wide region of parameter space. We computed optical CMDs for the donor stars and for the binary systems including radiation from the accretion disk. We also computed population diagrams showing the evolution of L_{pot} with donor age, and those showing the evolution of L_{pot} with P_{orb} .

When we plot the observed colors and magnitudes for six ULX systems found in the literature on the model CMDs, we find that all the data points lie in or near the high-probability regions spanned by the IMBH models on the CMD. On the other hand, fewer than half of the observed systems lie close to the high-probability region spanned by the LMBH models La and Lb, respectively. In light of this observation, we conclude that IMBH models are somewhat favored by the optical observations. The locations of the data on the CMDs indicate that these systems correspond to high mass donors, with $M_{\text{don}} \gtrsim 25 M_{\odot}$. This provides an indication as to why the IMBH models are favored over LMBH models. Our BPS calculations for LMBH binaries do not produce very many successful systems with a donor mass of $\gtrsim 20 M_{\odot}$, owing to limitations on the mass of the BH (~ 6 – $15 M_{\odot}$) and the requirement of stable mass transfer. On the other hand, the mass of the BH accretor in the IMBH models is set at $1000 M_{\odot}$, allowing for very massive donor stars, and hence the ability to account for higher optical luminosities.

We also find from the CMDs that the regions of high probability correspond to the initial phases of evolution of the donors. This is apparent because all stars spend a predominant fraction of their lifetimes on the main sequence. However, this is particularly consistent with the color-magnitude observations, all of which lie in or very near the high probability regions of the IMBH models. This leads us to conclude that the donor stars need to be predominantly on the main sequence or on the sub-giant branch. The $B-V$ color range of about -0.35 to -0.10 , corresponding to the high probability regions on the CMDs, indicates that the effective spectral class of the donors should be O through late B. We also dis-

cuss the effects of X-ray irradiation of the accretion disk, as well as intrinsic radiation generated viscously in the disk, on the different models. Disk radiation is found to be significant in the IMBH models, but less so in the LMBH models. We suggest that the main effects operating here are: (i) the higher donor masses in the IMBH systems, and hence higher values of L_{pot} , and (ii) the larger gravitational potential energy released at a given radial distance in the accretion disk in the IMBH models.

We compute population diagrams showing the evolution of L_{pot} with P_{orb} . As concluded previously from the CMDs, the $P_{\text{orb}}-L_{\text{pot}}$ diagrams show that the giant branch phases of evolution in these systems form a low-probability region in parameter space. It is seen that the most probable periods for all the models lie in the range of $\sim 1-10$ days, corresponding to the main-sequence and sub-giant phases of evolution. We quantify the likelihood of ULXs being in the pre-giant phase by estimating the relative probabilities for the systems to be in a pre-giant phase versus the giant phase. We find the probabilities to be in the ratios 0.08, 0.05 and 0.14 for Models La, Lb and Ic, respectively. We subsequently discuss the utility of the $P_{\text{orb}}-L_{\text{pot}}$ diagrams for estimating the probable range of P_{orb} for a system with a known L_{pot} .

Finally, we compute population diagrams showing the evolution of L_{pot} with the evolution time of the host star cluster. Similar diagrams have been reported for various ULX models in our previous studies (see Rappaport et al. 2005; Madhusudhan et al. 2006). In the present paper, we use the diagrams to calculate the ULX production efficiency as a function of the age of a star cluster, for the different models. We also estimate the numbers of ULXs in steady state for a typical galaxy as a function of L_{pot} . We find, with crude estimates, that the IMBH models explain the observed numbers of ULXs reasonably well, with ~ 0.07 with $L_{\text{pot}} > 10^{39}$ ergs s^{-1} per Milky Way type galaxy to ~ 0.03 per galaxy with $L_{\text{pot}} > 10^{40}$ ergs s^{-1} . For the LMBH models, the numbers range from about $\sim 12-20$ for $L_{\text{pot}} > 2 \times 10^{39}$ ergs s^{-1} per galaxy to $\sim 1.4-2.7$ with $L_{\text{pot}} > 10^{40}$ ergs s^{-1} . As discussed in §3.8 these numbers are too large by about a factor of 100 compared with the observations (Ptak & Colbert 2004; Grimm et al. 2003).

There are at least three ways of reducing the computed numbers of ULX systems in the LMBH model. One straightforward way is to adjust the λ parameter downward (see eq. [4] for a definition). We find that a value of λ in the range 0.01–0.03 will decrease the rate of production of BH binaries by the correct factors. Such small values of λ are quite consistent with calculations of envelope binding energies of massive stars (see Fig. 1 of Podsiadlowski et al. 2003; Dewi & Tauris 2000). Another possible explanation for the apparent overproduction of very luminous X-ray sources in the case of stellar-mass black-hole accretors is that the X-ray luminosities are, in fact, constrained to the Eddington limit, and that the large amounts of radiation-pressure ejected material severely attenuate the soft X-rays coming from the vicinity of the black hole. Finally, we mention the possibility that for case B mass transfer (i.e., when the primary progenitors of the black holes are in the Hertzsprung gap during the time when mass transfer commences) the primary is more likely to produce a neutron star rather than a black hole even for fairly high initial masses (e.g., $50-60 M_{\odot}$). As has first been argued by Brown et al.

(1999) and then has been confirmed in detailed calculations by Brown et al. (2001), this has to do with differences during helium core burning. Basically the final structure of a star is very different depending on whether the star burns helium with a hydrogen-burning shell around it or without the hydrogen-burning shell (as one would expect if the star loses its envelope before or early during helium burning). Without the hydrogen-burning shell the star ends up with a much smaller iron core at the end and most likely results in the production of a neutron star. This may be another reason why we do not find LMBH ULXs in nature.

In estimating the numbers of ULXs, we have allowed for violation of the Eddington limit by using L_{pot} in our calculations. This relaxation is imperative in order to explain ULXs using the LMBH models. However, the scenario by which the Eddington limit could be violated in a BH binary is not clear. We note that the numbers for the LMBH models with $L_{\text{pot}} > 10^{40}$ ergs s^{-1} could be quite unphysical because of the fact that for these luminosities $L_{\text{pot}} \gg L_{\text{Edd}}$. The Eddington limit for a $20 M_{\odot}$ accretor is $\sim 2.5 \times 10^{39}$ ergs s^{-1} . Allowing for the fact that a massive donor star could be transferring helium onto the accretor during the later stages of its evolution, the Eddington limit is increased by a factor of 2, to $\sim 5 \times 10^{39}$ ergs s^{-1} . Considering, however, that the bolometric luminosity may well be a factor of $\sim 2-3$ times the X-ray luminosities observed in the 1–10 keV X-ray band, the effective Eddington limit in the X-ray band is only $\sim 1.5 \times 10^{39}$ ergs s^{-1} . Thus, even if we allow for a violation of the Eddington limit by reasonable factors of a few, and thereby achieve L_{pot} up to $\sim 10^{40}$ ergs s^{-1} , it is difficult to understand X-ray-band luminosities above $\sim 5 \times 10^{39}$ ergs s^{-1} .

As for the IMBH model, we have not studied in any detail the formation scenarios of IMBHs or of the capture of binary companions by the IMBHs (but see Portegies Zwart et al. 2004; Tutukov & Fedorova 2005; Blecha et al. 2006). Despite much evidence in the literature supporting IMBH binaries as candidates for ULXs, the formation mechanisms of IMBHs are not clear. In particular, it is unclear how supermassive stars evolve, and whether they can undergo core collapse to form an IMBH. In calculating the numbers of ULXs in this scenario, we have assumed plausible values for the formation rate of young star clusters in a galaxy, the fraction of these clusters that successfully produce IMBHs, and the probability of capture of a massive companion by the IMBH.

Finally, we should keep in mind the possibility that neither the IMBH nor LMBH binary models may be correct in explaining the most luminous ULXs. It is conceivable that the most luminous ULXs may, in fact, be free floating IMBHs that are fed by tidally disrupted passing field stars (see Volonteri & Perna 2005, for a related scenario).

We are grateful to Bill Paxton for all his assistance with the EZ stellar evolution code. We thank Wlodek Kluźniak for helpful discussions regarding accretion disks. SR acknowledges support from NASA Chandra Grant TM5-6003X. LN thanks NSERC (Canada) and the CRC Program for financial support, and also acknowledges the CCS at the Universite de Sherbrooke for their technical assistance.

REFERENCES

Abramowicz, M.A., Czerny, B., Lasota, J.P., & Szuszkiewicz, E. 1988, *ApJ*, 332, 646.

Bardeen, J.M. 1970, *Nature*, 226, 64.

- Begelman, M. 2002, *ApJ*, 568, 97.
- Begelman, M. 2006, *ApJ*, 643, 1065.
- Blecha, L., Ivanova, N., Kalogera, V., Belczynski, K., Fregeau, J., & Rasio, F. 2006, *ApJ*, 642, 1.
- Brown, G. E., Lee, C.-H., & Bethe, H. A. 1999, *NewA*, 4, 313.
- Brown, G. E., Heger, A., Langer, N., Lee, C.-H., Wellstein, S., & Bethe, H. A. 2001, *NewA*, 6, 457.
- Colbert, E., & Mushotzky, R. 1999, *ApJ*, 519, 89.
- Cropper, M., Soria, R., Mushotzky, R.F., Wu, K., Markwardt, C.B., & Pakull, M. 2004, *MNRAS*, 349, 39.
- Copperwheat, C., Cropper, M., Soria, R., Wu, K. 2005, *MNRAS*, 362, 79.
- Copperwheat, C., Cropper, M., Soria, R., Wu, K. 2007, *MNRAS*, 376, 1407.
- Dewi, J. & Tauris, T. 2000, *A&A*, 360, 1043.
- de Grijs, R., Anders, P., Bastian, N., Lynds, R., Lamers, H.J.G.L.M., O’Neil, Jr, E. J. 2003, *MNRAS*, 343, 1285.
- Ebisawa, K., Życki, P., Kubota, A., Mizuno, T., Mushotzky, R., Watarai, K. Y. 2003, *ApJ*, 597, 780.
- Eggleton, P.P. 1983, *ApJ*, 268, 368.
- Grimm, H.-J., Gilfanov, M., Sunyaev, R. 2003, *MNRAS*, 339, 793.
- Hopman, C., Portegies Zwart, S.F., & Alexander, T. 2004, *ApJ*, 604, L101.
- Hunter, D. A., Elmegreen, B. G., Dupuy, T. J., Mortonson, M. 2003, *ApJ*, 126, 1836.
- Jaroszynski, M., Abramowicz, M. A., Paczyński, B. 1980, *Acta Astronomica*, Vol. 30.1.
- Johnson, H. L. 1966, *ARA&A*, 4, 193.
- Kaaret, P., Ward, M.J., & Zezas, A. 2004, *MNRAS*, 351, L83.
- King, A.R., Kolb, U., Szuszkiewicz, E. 1997, *ApJ*, 488, 89.
- King, A.R., Davies, M.B., Ward, M.J., Fabbiano, G., & Elvis, M. 2001, *ApJ*, 552, L109.
- Körding, E., Falcke, H., & Markoff, S. 2002, *A&A*, 382, L13.
- Kuntz, K. D., Gruendl, R. A., Chu, Y.-H., Chen, C.-H. Rosie., Still, M., Mukai, K., Mushotzky, R. F. 2005, *ApJ*, 620, L31.
- Liu, J.-F., Bregman, J. N., Seitzer, P. 2002, *ApJ*, 580, L31.
- Madhusudhan, N., Justham, S., Nelson, L., Paxton, B., Pfahl, E., Podsiadlowski, Ph., Rappaport, S. 2006, *ApJ*, 640, 918.
- Miller, J.M., Fabbiano, G., Miller, M.C., & Fabian, A.C. 2003, *ApJ*, 585, L37.
- Miller, J.M., Fabian, A.C., & Miller, M.C. 2004, *ApJ*, 607, 931.
- Mucciarelli, P., Zampieri, L., Falomo, R., Turolla, R., Treves, A. 2005, *ApJ*, 633, L101.
- Mucciarelli, P., Zampieri, L., Treves, A., Turolla, R., Falomo, R. 2007, *ApJ*, 658, 999.
- Pakull, M., & Mirioni, L. 2003, *RMxAC*, 15, 197.
- Paxton, B. 2004, *PASP*, 116, 699.
- Pfahl, E., Rappaport, S., & Podsiadlowski, Ph. 2003, *ApJ*, 597, 1036.
- Pfahl, E. 2005, *ApJ*, 626, 849.
- Podsiadlowski, Ph., Rappaport, S., & Han, Z. 2003, *MNRAS*, 341, 385.
- Pols, O. R., 1994, *A&A*, 290, 119.
- Pols, O.R., Tout, C.A., Eggleton, P.P. & Han, Zh. 1995, *MNRAS*, 274, 964.
- Pooley, D., Rappaport, S. 2005, *ApJ*, 634, L85.
- Portegies Zwart S.F., Dewi, J., & Maccarone, T. 2004, *MNRAS*, 355, 413.
- Ptak, A., & Colbert, E. 2004, *ApJ*, 606, 291.
- Rappaport, S., Podsiadlowski, Ph., & Pfahl, E. 2005, *MNRAS*, 356, 401.
- Shakura, N. I., Sunyaev, R. A. 1973, *A&A*, 24, 337.
- Socrates, A., & Davis, S. W. 2006, *ApJ*, 651, 1049.
- Soria, R., Cropper, M., Pakull, M., Mushotzky, R., Wu, K. 2005, *MNRAS*, 356, 12.
- Strohmayer, T.E., & Mushotzky, R.F. 2003, *ApJ*, 586, 61.
- Tutukov, A.V., & Fedorova, A.V. 2005, *Astronomy Reports*, 49, 89.
- Volonteri, M., & Perna, R. 2005, *MNRAS*, 358, 913.
- Webbink, R. 1985, in *Interacting Binary Stars*, ed. J.E. Pringle & R.A. Wade (Cambridge: Cambridge Univ. Press), 39.
- Wellstein, S., Langer, N., & Braun, H. 2001, *A&A*, 369, 939.